

Last time: More on Stokes' theorem

Consider the complicated surface S drawn on the board. Let $\mathbf{F} = \langle P, Q, R \rangle$ be a vector field (with continuous first order partial derivatives) defined on an open set D containing S . What can you say about $\iint_S \text{curl} \mathbf{F} \cdot d\mathbf{S}$?

- (a) It's zero, because ∂S is empty.
- (b) It's not defined unless \mathbf{F} is defined over the entire solid bounded by the surface S .
- (c) We can't say anything unless we know more about \mathbf{F} .
- (d) I don't know.

If you want to come to extra office hours/review session, fill out the form on the course diary. You can do it right now, if you're done!

Physical meaning of div

Recall that for a fluid flow \mathbf{F} , the **flux** of \mathbf{F} across an oriented surface S measures the amount of fluid crossing S (in the direction of the positive normal vector) in unit time. It is calculated by

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_S \mathbf{F} \cdot \mathbf{n} dS.$$

So if S is the boundary of some solid E (oriented to point away from E), the Divergence Theorem tells us that

$$\text{Flux} = \iint_{\partial E} \mathbf{F} \cdot d\mathbf{S} = \iiint_E \text{div} \mathbf{F} \, dV.$$

Physical meaning of div

So

$$\text{Flux} = \iiint_E \text{div}\mathbf{F} \, dV.$$

If E is a tiny ball with volume $V(E)$ centered around a point P , we approximate the flux as follows:

$$\begin{aligned} \iiint_E \text{div}\mathbf{F} \, dV &= V(E) \cdot (\text{average value of } \text{div}\mathbf{F} \text{ on } E) \\ &\approx V(E) \cdot \text{div}\mathbf{F}(P). \end{aligned}$$

So

- Fluid is **leaving** E when $\text{div}\mathbf{F}(P) > 0$ —we say P is a **source**
- Fluid is **entering** E when $\text{div}\mathbf{F}(P) < 0$ —we say P is a **sink**
- If $\text{div}\mathbf{F}(P) = 0$, the total amount of fluid leaving E is equal to the total amount of fluid entering E .

Calculating flux using the divergence theorem

Given E , S and S' as on the board, what is

$$\iint_S \mathbf{F} \cdot d\mathbf{S}?$$

- (a) 16
- (b) 4
- (c) -4
- (d) -12
- (e) I don't know.

Summary: We wanted to find the integral of \mathbf{F} over S . S was pretty hard, because it has five faces.

But we could put a lid S' on S , making it into the boundary of a solid box E .

- So $\iint_S \mathbf{F} \cdot d\mathbf{S} + \iint_{S'} \mathbf{F} \cdot d\mathbf{S} = \iint_{\partial E} \mathbf{F} \cdot d\mathbf{S}$.
- $\iint_{S'} \mathbf{F} \cdot d\mathbf{S}$ is pretty easy to compute.
- $\iint_{\partial E} \mathbf{F} \cdot d\mathbf{S}$ would be hard to compute directly (because it has **six** faces!), but it's the boundary of a solid E , so we have a trick—the divergence theorem tells us that

$$\iint_{\partial E} \mathbf{F} \cdot d\mathbf{S} = \iiint_E \operatorname{div} \mathbf{F} \, dV.$$

Example

Let $S = \{(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2\}$, ($r > 0$). Let $F = \langle x, y, z \rangle$. How much fluid flows across S in unit time?

- (a) πr^3
- (b) $4\pi r^3$
- (c) $\frac{4}{3}\pi r^3$
- (d) The answer depends on (x_0, y_0, z_0) and r .
- (e) The calculation is too complicated.