

## Last time: the Divergence Theorem

Assume  $\mathbf{E}$  is a vector field with continuous first derivatives on an open set  $D \subset \mathbb{R}^3$ . Assume that  $B \subset D$  is a “nice” solid. Then

### Theorem (Divergence Theorem)

$$\iiint_B \operatorname{div} \mathbf{E} \, dV = \iint_{\partial B} \mathbf{E} \cdot d\mathbf{S}.$$

**Example.** Assume  $D = \mathbb{R}^3 \setminus \{(0, 0, 0)\}$  is everything except the origin. Suppose that  $\operatorname{div} \mathbf{E} = 0$  on  $D$ . Let  $S_r = \{x^2 + y^2 + z^2 = r^2\}$  and let  $S'_r = \{(x - 3)^2 + (y - 1)^2 + z^2 = r^2\}$ .

Find  $I_1 = \iint_{S_1} \mathbf{E} \cdot d\mathbf{S}$  and  $I_2 = \iint_{S'_1} \mathbf{E} \cdot d\mathbf{S}$ .

- (a) Not enough information to find either.
- (b)  $I_1 = I_2 = 0$ .
- (c) Not enough information to find  $I_1$ ; but  $I_2 = 0$ .
- (d)  $I_1 = 0$ ; not enough information to find  $I_2$ .

## Announcements

- Final exam is next Friday. Register for conflict by Monday.
- Office hours/review session next week:
  - Ordinary office hours Tuesday 11–11:50am.
  - Extra office hours Wednesday evening (probably 6–7pm, maybe 7–8pm—it's fine with me if you bring your dinner). **AH 341**
  - Extra office hours Thursday 12–1pm. **AH 341**
  - Possibly office hours also on Friday, but I can't confirm yet.
  - Come with questions (or you can listen to other people's questions).
- Fact: today is our ante-penultimate lecture. Wednesday was our **pre-ante-penultimate** lecture, but I forgot to say so.

## Electric field and electric flux

Given a particle of charge  $Q$  at  $(0, 0, 0)$ , its **electric field** is

$$\mathbf{E}(x, y, z) = \frac{Q}{4\pi\epsilon_0(x^2 + y^2 + z^2)^{\frac{3}{2}}} \langle x, y, z \rangle$$

or equivalently

$$\mathbf{E}(\mathbf{r}) = \frac{Q}{4\pi\epsilon_0|\mathbf{r}|^3} \mathbf{r}. \quad \text{Inverse square law}$$

This means that the force experienced by a particle of charge  $q$  at position  $\mathbf{r}$  is  $q\mathbf{E}(\mathbf{r})$ .

Where is the vector field  $\mathbf{E}$  defined?

- (a) all of  $\mathbb{R}^3$
- (b) everywhere except  $(0, 0, 0)$
- (c) everywhere except the  $z$ -axis,  $\{x = y = 0\}$
- (d) I don't know

## Practice with Gauss' Law

Suppose we have particles of charge  $Q_i$  at points  $P_i$ , with  $Q_i = i$  for  $i = 1, 2, 3, 4, 5$ . Suppose that  $B$  is a solid region containing  $P_1$ ,  $P_3$ , and  $P_4$ , but not  $P_2$  or  $P_5$ . What is

$$\iint_{\partial B} \mathbf{E} \cdot d\mathbf{S}?$$

- (a) 0
- (b)  $\frac{1}{\epsilon_0}$
- (c)  $\frac{2}{\epsilon_0}$
- (d)  $\frac{4}{\epsilon_0}$
- (e)  $\frac{8}{\epsilon_0}$